

- 3 **Lesperance F**, Frasere-Smith N, Juneau M, *et al.* Depression and 1-year prognosis in unstable angina. *Arch Intern Med* 2000;**160**:1354–60.
- 4 **Lesperance F**, Frasere-Smith N, Talajic M, *et al.* Five-year risk of cardiac mortality in relation to initial severity and one-year changes in depression symptoms after myocardial infarction. *Circulation* 2002;**105**:1049–53.
- 5 **Bush DE**, Ziegelstein RC, Tayback M, *et al.* Even minimal symptoms of depression increase mortality risk after acute myocardial infarction. *Am J Cardiol* 2001;**88**:337–41.
- 6 **Steeds RP**, Channer KS. Depression: the sleeping giant. *Eur Heart J* 2000;**21**:427–9.
- 7 **Frasere-Smith N**, Lesperance F. Reflections on depression as a cardiac risk factor. *Psychosom Med* 2005;**67**(suppl 1):S19–25.
- 8 **Lane D**, Ring C, Lip GY, *et al.* Depression, indirect clinical markers of cardiac disease severity, and mortality following myocardial infarction. *Heart* 2005;**91**:531–2.
- 9 **Lane D**, Carroll D, Lip GY. Anxiety, depression, and prognosis after myocardial infarction: is there a causal association? *J Am Coll Cardiol* 2003;**42**:1808–10.
- 10 **Shimbo D**, Davidson KW, Haas DC, *et al.* Negative impact of depression on outcomes in patients with coronary artery disease: mechanisms, treatment considerations, and future directions. *J Thromb Haemost* 2005;**3**:897–908.
- 11 **Glassman AH**, O'Connor CM, Califf RM, *et al.* Sertraline treatment of major depression in patients with acute MI or unstable angina. *JAMA* 2002;**288**:701–9.
- 12 **Berkman LF**, Blumenthal J, Burg M, *et al.* Effects of treating depression and low perceived social support on clinical events after myocardial infarction: the enhancing recovery in coronary heart disease patients (ENRICH) randomized trial. *JAMA* 2003;**289**:3106–16.
- 13 **Swenson JR**. Quality of life in patients with coronary artery disease and the impact of depression. *Curr Psychiatry Rep* 2004;**6**:438–45.
- 14 **Wells KB**, Stewart A, Hays RD, *et al.* The functioning and well-being of depressed patients. Results from the medical outcomes study. *JAMA* 1989;**262**:914–9.
- 15 **Spertus JA**, Jones P, McDonell M, *et al.* Health status predicts long-term outcome in outpatients with coronary disease. *Circulation* 2002;**106**:43–9.
- 16 **Ruo B**, Rumsfeld JS, Hlatky MA, *et al.* Depressive symptoms and health-related quality of life: the heart and soul study. *JAMA* 2003;**290**:215–21.
- 17 **Schleifer SJ**, Macari-Hinson MM, Coyle DA, *et al.* The nature and course of depression following myocardial infarction. *Arch Intern Med* 1989;**149**:1785–9.
- 18 **Lesperance F**, Frasere-Smith N, Talajic M. Major depression before and after myocardial infarction: its nature and consequences. *Psychosom Med* 1996;**58**:99–110.
- 19 **Spertus JA**, McDonell M, Woodman CL, *et al.* Association between depression and worse disease-specific functional status in outpatients with coronary artery disease. *Am Heart J* 2000;**140**:105–10.
- 20 **Lane D**, Carroll D, Ring C, *et al.* Mortality and quality of life 12 months after myocardial infarction: effects of depression and anxiety. *Psychosom Med* 2001;**63**:221–30.
- 21 **de Jonge P**, Spijkerman TA, van den Brink RHS, *et al.* Depression after myocardial infarction is a risk factor for declining health related quality of life and increased disability and cardiac complaints at 12 months. *Heart* 2006;**92**:32–9.
- 22 **Swenson JR**, O'Connor CM, Barton D, *et al.* Influence of depression and effect of treatment with sertraline on quality of life after hospitalization for acute coronary syndrome. *Am J Cardiol* 2003;**92**:1271–6.
- 23 **Ziegelstein RC**, Kim SY, Kao D, *et al.* Can doctors and nurses recognize depression in patients hospitalized with an acute myocardial infarction in the absence of formal screening? *Psychosom Med* 2005;**67**:393–7.

IMAGES IN CARDIOLOGY


doi: 10.1136/hrt.2005.063800

An unexpected complication during right jugular vein cannulation in a young adult

The figure presents a complete atrioventricular (AV) block which developed in a 35 year old woman hospitalised in our intensive care unit because of septic shock. She had been first hospitalised in the gastroenterology department because of an upper gastrointestinal tract haemorrhage caused by oesophageal varices in the context of hepatic cirrhosis of unknown cause. The AV block developed when we attempted to insert a right jugular vein catheter several days later. During the insertion of the guidewire the patient developed a brief run of atrial tachycardia followed by a period of complete atrioventricular block. Her arterial pressure fell dramatically. The guidewire was partially withdrawn and her cardiac rhythm and pressure were restored to normal.

Our patient did not present any cardiac abnormality. The admission ECG demonstrated a sinus tachycardia (136 beats/min) and a P–R interval of 0.12 s while cardiac enzyme measurements performed the days that followed her admission were normal. After the event, factors that could explain the block such as acidosis, electrolyte disturbances, endocarditis, large ascitic fluid quantity, and medications (for example, β blockers) were ruled out.

Arrhythmias during central venous cannulation are more prone to occur with the insertion of a pulmonary arterial catheter, especially if a pre-existing left bundle branch block (LBBB) is present. In literature, the induction of a complete heart block during the insertion of a central venous catheter has been described in two older patients with a pre-existing LBBB. The most probable explanation in our patient is a transient injury to the distal AV node caused by the over-insertion of the guide wire. Care must be taken when inserting guidewires under continuous monitoring in a distance less than 16–18 cm, since life threatening arrhythmias can occur even in young patients without cardiac abnormalities.



E Paramythiotou
I Poularas
A Kalogeromitros
V Krikos
N Katsarelis
A Karabinis

lparamyth61@hotmail.com